

Balance D

Contains ingredients important for the synthesis of dopamine, a catecholamine important for positive affect, optimal mood, cognition, and craving control*


Item Number	Available Sizes	Serving Size
2067	60 Capsules	2 Capsules



Key Ingredients

N-acetyl-L-cysteine	<ul style="list-style-type: none"> ■ Precursor to L-cysteine¹ ■ L-cysteine is required for the synthesis of glutathione, an endogenous antioxidant²
N-acetyl-L-tyrosine	<ul style="list-style-type: none"> ■ L-tyrosine is a precursor to catecholamines including dopamine, norepinephrine, and epinephrine³
Mucuna cochinchinensis seed extract (99% L-DOPA)	<ul style="list-style-type: none"> ■ Natural source of L-DOPA⁴ ■ L-DOPA crosses the blood-brain barrier and is the direct precursor to dopamine⁵
Vitamins B6, C, and folate	<ul style="list-style-type: none"> ■ Active forms of vitamin B6, C, and folate are important for catecholamine synthesis^{6-8*}

The Science




Catecholamines

A class of neurotransmitters responsible for many functions in the nervous and endocrine systems⁹

- Catecholamines play an important role in optimal **mood**, energy, **memory**, attention, and **cognition**¹⁰⁻¹³


Catecholamine Pathway



*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

MORE SCIENCE BEHIND BALANCE D

Figure 1. Dopaminergic Signaling in the Brain



Dopamine is important for craving control


Dopaminergic function is critical in balancing cravings, reward based behaviors, and healthy weight management¹⁴

- Dopamine release from the ventral tegmental area (VTA) to the nucleus accumbans is the primary pathway in which reward based behaviors such as cravings are modulated¹⁵
- Impaired signaling from decreased dopamine release or receptors (DRD2) is thought to reduce sensitivity to natural reward-based behavior¹⁶

Balance D provides ingredients to support dopamine synthesis^{3,6*}

- N-acetyl-L-tyrosine and *Mucuna cochinchinensis* seed extract (99% L-DOPA) are amino acid precursors to dopamine³
- Vitamin B6 is a necessary cofactor for the synthesis of dopamine^{6*}

Figure 2. Oxidative Stress Affects Neuronal Health




Formulated for a comprehensive pathway approach

Produced in excess, dopamine is oxidized into quinones, free radicals, and reactive oxygen species (ROS)¹⁷


- Quinones react with glutathione (GSH) and deplete levels of the reduced GSH and total antioxidant capacity¹⁸
- Excessive production of ROS and decreased antioxidant function contributes to cellular damage, dysfunction, and/or death¹⁷

Balance D contains ingredients for the synthesis of glutathione

- N-acetyl-L-cysteine is the precursor to L-cysteine; L-cysteine is an amino acid precursor in the synthesis of glutathione^{1,2}
- Selenium is a micronutrient that helps to comprise selenoproteins; these proteins are crucial for brain function and the enzyme glutathione peroxidase (GPx)^{19*}



1. Rushworth G and Megson I. *Pharmacol Ther.* 2014;141(2):150-9.
2. Schmitt B, et al. *Redox Biol.* 2015;6:198-205.
3. Ng J, et al. *Nat Rev Neurol.* 2015;11:567-84.
4. Tuleun C, et al. *Livestock Research for Rural Development.* 2008;20(10).
5. Partridge W. *NeuroRx.* 2005;2(1):3-14.
6. Dakshinamurti K, et al. *Ann N Y Acad Sci.* 1990;585:128-44.
7. Levine M, et al. *J Biol Chem.* 1985;260(24):12942-7.
8. Stahl S. *J Clin Psychiatry.* 2008;69(9):1352-3.
9. Eisenhofer G, et al. *Pharmacol Rev.* 2004;56(3):331-49.
10. Clark K and Noudoost B. *Front Neural Circuits.* 2014;8:33.



Add more mood support?

Learn more about TravaCor at www.neuroscienceinc.com/products/travacor

11. Blier P. *J Psychiatry Neurosci.* 2001;26 Suppl:S1-2.
12. Verhoeff N, et al. *Pharmacol Biochem Behav.* 2003;74(2):425-32.
13. Xing B, et al. *Brain Res.* 2016;1641(Pt B):217-33.
14. Blum K, et al. *Front Psychol.* 2014;5:919.
15. Wise R. *Philos Trans R Soc Lond B Biol.* 2006;361(1471):1149-58.
16. Geiger BM, et al. *FASEB J.* 2008;22(8):2740-6.
17. Smeysne M and Smeysne R. *Free Radic Biol Med.* 2013;62:13-25.
18. Spencer J, et al. *J Neurochem.* 1998;71(5):2112-22.
19. Rayman M. *Lancet.* 2012;379:1256-69.

*Magnafolate is a registered trademark of Lianyungang Jinkang Pharmaceutical Technology Co., Ltd.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

(888) 342-7272 www.neuroscienceinc.com

Copyright © 2024 NeuroScience 031824-N2103